

Entropy Defined, Entropy Increase and Decoherence Understood, and Some Black-Hole Puzzles Solved

Bernard S. Kay, Department of Mathematics, University of York, York YO1 5DD, UK

Statistical mechanics explains thermodynamics in terms of (quantum) mechanics by equating the entropy of a microstate of a closed system with the logarithm of the number of microstates in the macrostate to which it belongs, but the question ‘what is a macrostate?’ has never been answered except in a vague, subjective, way. However Hawking’s discovery^{1,2} of black hole evaporation led to a formula for black hole entropy with no subjective element. In this letter, we argue from this result, together with the assumption³ that ‘black hole thermodynamics is just ordinary thermodynamics applied to black holes’, that a macrostate for a general (quantum gravitational) closed system is *an equivalence class of matter-gravity microstates with the same expectation values for the matter degrees of freedom alone*. Not only does this finally answer the question ‘what is entropy?’, but it also predicts the equality of the thermodynamic entropy of a black hole with the matter and the gravity entropy-like quantities derived^{4,5} from the Euclidean path integral. Furthermore it gives us a clear glimpse of an ultimate synthesis of quantum theory and gravity in which we see that (a) gravity acts as a *universal environment*, thus predicting that, if the initial state of the universe is unentangled, its entropy must go on increasing forever, (b) *the gravitational field has degrees of freedom, but no observables*, thus enabling gravity to perform the trick of providing *an objective continual process of decoherence*. All the above rests on the validity of unitarity. The ‘information-loss puzzle’⁶ had raised doubts about that. But we suggest a resolution for this puzzle.

Our thesis is that thermodynamics can be derived as a consequence of the microscopic laws of nature only when these microscopic laws are taken to be the laws of quantum gravity. Relying only on basic, conservative, assumptions about the quantum mechanical framework underlying quantum gravity, together with well-established results from the subject of (quantum) ‘black-hole thermodynamics’, not only do we then find that entropy acquires an objective definition, but also a new paradigm for the laws of nature emerges which seems capable of solving many of the longstanding puzzles of theoretical physics and cosmology. Moreover, the new paradigm corrects our previous understanding of black hole thermodynamics in a way which explains some previously unexplained coincidences.

We must first clarify what we mean by a closed system in quantum gravity. Because of interactions, and because gravity is ‘universal’, i.e. it couples to all matter, this can only be a model which either is intended literally to be a description of the whole universe, or which describes our system of interest *as if* it were the whole universe. For example, if one were interested in the physics of a single neutron star, one would consider a fictitious universe consisting of a single neutron star sitting in an empty, apart from quantum vacuum fluctuations, asymptotically flat space. Fictitious systems in ‘boxes’ are also permitted, with due regard to appropriate fictitious boundary conditions, in the same spirit.

Our starting point is to posit the existence, and correctness, of a quantum gravity theory which is in conformity with standard quantum mechanics in the sense that the following two assumptions hold: First, the theory assigns independent dynamical degrees of freedom for gravity and matter so the total Hilbert space for a closed system will take the, tensor product, form

$$H = H_{\text{matter}} \otimes H_{\text{gravity}}.$$

Second, any full description of any closed system corresponds to the standard quantum mechanical notion of a *single microstate* – i.e. to a density operator ρ which takes the form $|\Psi\rangle\langle\Psi|$ where $|\Psi\rangle$ is a vector, or ‘total wave function’ in H ; in other words, to what is usually called a ‘pure total state’. We shall postpone the question of ‘what is time’ to after our discussion of entropy increase below, and, as an interim interpretation, simply think of the total Hilbert space as representing the possible microstates of the system ‘at a given single time’.

By standard quantum mechanics⁷, we know that, given any total Hilbert space H which takes the form $H_a \otimes H_b$, (we are interested in the case where a stands for *matter* and b for *gravity*) and a microstate $\rho = |\Psi\rangle\langle\Psi|$, then, if we are only concerned about an observable A tied to the a degrees of freedom, the expectation value $\langle A \rangle$ of A is given by a density operator ρ_a on H_a , namely the ‘partial trace’ of ρ over the Hilbert space H_b , satisfying

$$\text{trace}(\rho A \otimes I_b)_H = \text{trace}(\rho_a A)_{H_a} \quad (1)$$

where I_b stands for the identity operator on H_b . Similarly there will be ρ_b on H_b , suitable for observables B tied to the b degrees of freedom, and by the standard von Neumann procedure⁷, we can assign to each of these partial states the ‘partial entropies’ $S_a = -\text{trace}(\rho_a \log \rho_a)$ and $S_b = -\text{trace}(\rho_b \log \rho_b)$, which, as we explain below, and as is well known, must (as a consequence of the purity of the total state) necessarily be equal (and, of course, non-negative). Of course, the total von Neumann entropy of the microstate ρ , will be zero, and when we consider a time-evolution, below, will remain zero for all time. Thus ‘additivity of the entropy’ fails to the maximum possible extent here. *But in the sequel, the von Neumann entropy of the total state will have no direct physical interpretation.* ρ_a is the proper mathematical representation of what we call here an ‘*a macrostate*’. In the case ρ_a is a projector onto an n -dimensional subspace, then one may identify the ‘*number of microstates in the macrostate ρ_a* ’ with the number n . But, we shall continue to use this phrase, picturesquely, also to describe the general case. This way we can think of the a macrostates as *the equivalence classes of microstates which are indistinguishable from one another, as far as the a observables are concerned*. The partial entropy S_a may then be thought of as the logarithm of the number of microstates for the total system contained in the macrostate ρ_a . A corresponding statement holds for the interpretation of S_b ($= S_a$).

It will be helpful to recall how the equality $S_a = S_b$ may be proven: The total wave function $|\Psi\rangle$ may be written as a superposition, with coefficients c_i satisfying $\sum_i |c_i|^2 = 1$, of single tensor products

$$|\Psi\rangle = \sum_i c_i |\alpha_i\rangle \otimes |\beta_i\rangle$$

where the $|\alpha_i\rangle$ and $|\beta_i\rangle$ (which depend on $|\Psi\rangle$) are orthonormal sets of basis vectors in H_a and H_b respectively. One easily has from this that

$$\rho_a = \sum_i |c_i|^2 |\alpha_i\rangle\langle\alpha_i| \quad \text{and} \quad \rho_b = \sum_i |c_i|^2 |\beta_i\rangle\langle\beta_i|$$

and from this it easily follows that

$$S_a = S_b = - \sum_i |c_i|^2 \log |c_i|^2.$$

Finally, we remark that when the total wave function can be written as a single non-zero tensor product, the total state is often said to be unentangled. In this case, the partial entropies will be zero. For this reason, the value of the (equal) partial entropies in the general case, is often referred to as ‘entanglement entropy’.

Returning to the case of interest, where a stands for *matter* and b for *gravity*, we shall now argue that the value of the *physical entropy*, S , of a matter-gravity microstate $\rho = |\Psi\rangle\langle\Psi|$ must be the value of the (equal) partial matter and gravity entropies of ρ , i.e. it must be its ‘matter-gravity entanglement entropy’. In particular,

$$S = S_{\text{matter}} \quad (= -\text{trace}(\rho_{\text{matter}} \log \rho_{\text{matter}})) \quad (2)$$

or, in words:

The physical entropy, S , of the microstate ρ is the logarithm of the number of matter-gravity microstates for which the expectation values of all the matter degrees of freedom coincide with the expectation values in ρ .

More precisely we shall argue that, if there is any quantity at all in quantum gravity, which is a function on the set of all possible microstates, taking values in the non-negative real numbers, and which coincides with what we call ‘entropy’ both in every-day and in black-hole thermodynamics, then it must be given by (2).

We shall give two entirely independent arguments for this statement. We feel that, while each argument, by itself, might be open to some reasonable doubt, the two arguments taken together make a very strong case that (2) is true. Our first argument begins by observing that, in the thermodynamics of everyday closed systems, one is concerned with measurements of the matter, but not of the gravity, degrees of freedom. Thus the (vague, ill-defined) old statistical-mechanical entropy is equal to

- (I) the entropy one would obtain, by modifying the von Neumann formula with the usual ‘coarse graining procedures’, when one coarse-grains in a way appropriate to measurements of the matter degrees of freedom with a given degree of experimental accuracy, and gravity is not measured at all.

On the other hand, (2) asserts that the physical entropy is

- (II) the entropy obtained when one ‘coarse grains’ as if measurements of the matter degrees of freedom could be made (to within the fundamental limitations of the Heisenberg uncertainty principle) with *maximal* accuracy, and gravity is *still* not measured at all.

Because of the well-known robustness of the traditional arguments of statistical mechanics under changes in the notion of coarse graining, it seems plausible that, as long as there's a reasonable number of gravitational degrees of freedom, it will not make any difference which of these two coarse grainings we actually use. If it did, thermodynamics would start going wrong when one started to make measurements of matter too accurately. Thus, we conclude that in every-day thermodynamics, while we have a lot of choice as to how entropy may be defined, the entropy *may be taken to be* the S of (2), and this is the only definition which is free from any subjective element; *gravity provides an ideal, and objective, notion of coarse-graining*. But in the case of a *black hole* we combine our knowledge of Hawking's formula *temperature equals surface gravity over 2π* with the hypothesis³ that 'black hole thermodynamics is ordinary thermodynamics applied to black holes' to conclude that the entropy must equal

- (i) the entropy obtained using the thermodynamic formula $dE = TdS + \text{work terms}$, from the above temperature formula.

This takes the value² 'one quarter of the area of the event horizon' and the important point for our argument is that this is a definite value, free from any subjective element. We conclude that if we are to have a single formula valid for *all* closed systems, both every-day and black-hole, the physical entropy has to be given by (2).

Our second argument takes as its starting point the remarkable result⁴ from the 'Euclidean path-integral approach'⁵ to quantum black holes that another 'entropy-like quantity' namely

- (ii) the 'entropy' obtained^{4,5} from the pure gravity partition function^{4,5} for a quantum black hole

turns out, by a hitherto unexplained coincidence, to equal, to zero-loops^{4,5}, one quarter of the area of the event horizon, i.e. the same value as (i). On the other hand, we shall argue below that (ii) ought (to zero loops) to be equated with S_{gravity} which we know must equal S_{matter} . Thus we conclude that (2) is true for the case of a black hole. By considering examples of closed systems which include both black holes and every-day objects, one can further argue that this conclusion must extend to general microstates.

Once one accepts the definition (2) for physical entropy, the prospect opens up of being able, once we have a theory of quantum gravity, to replace the traditional, partially unsatisfactory, explanation of thermodynamics in terms of a (quantum) 'statistical mechanics', by an explanation in terms of a quantum-gravitational 'mechanics'. In fact, we can already see the qualitative shape such an explanation will take: The laws of thermodynamics should emerge for the *matter degrees of freedom* regarded as an *open system* in contact with the *environment* provided by the gravitational field, and coupled to the latter according to the interactions determined by the physically correct full theory of quantum gravity when the entropy is regarded as the attribute of each microstate of the total *matter-gravity closed system* defined by (2). The fact that the gravitational coupling is so weak, encourages us to expect that an explanation along these lines will be successful.

We have now arrived at the right place to discuss the, interrelated, problems of 'time' and 'entropy increase'. In order to proceed with the enterprise outlined above along anything like traditional lines, one will need a notion for a flow of time, represented by a

one-parameter family of unitary operators, which we shall call a *flow* $t \mapsto U(t)$ on H with t ranging over the non-negative real numbers, mapping any ‘initial microstate’ $\rho_0 = |\Psi_0\rangle\langle\Psi_0|$ at some ‘initial time’ $t = 0$ into the microstate $\rho_t = |\Psi_t\rangle\langle\Psi_t|$ at a ‘later time’, t , according to the transformation

$$\rho_t = U(t)\rho_0U(t)^{-1}. \quad (3)$$

So a single microstate at one time evolves to a single microstate at a later time. It is very important to realize that it will be the theory of quantum gravity which will, presumably, tell us which such flows deserve to be called *time-flows*. (In fact, it seems conceivable that a quantum theory of gravity *is* just a specification of which flows are time-flows.) The problem of the increase of entropy of a closed system seems now susceptible of solution, once we add the assumption, for any closed system of interest, that the initial matter-gravity microstate is totally unentangled. Its initial entropy will then obviously be zero, and it is obvious, that, at least at the start, its entropy $S(t)$ can only increase. But we should be able to say much more: There would seem to be very good hopes that arguments similar to the traditional arguments of statistical mechanics for open systems in contact with environments, but now of a deterministic kind, will, for the time-flows sanctioned by quantum gravity, now ensure that $S(t)$ must go on increasing for ever and can never turn around and start decreasing at some time. In particular, given such a notion of time-flow, if we assume the initial microstate ρ_0 , has zero, or low, entropy then one should be able to argue that *the entropy of the universe must go on increasing forever*. This way of explaining entropy increase is to be contrasted with attempts within the traditional paradigm. While something like the above picture can be argued for with a suitable notion of coarse-graining, as long as one believed that, ultimately, physical entropy should be identified with the von Neumann entropy of the total state, then, assuming the universe to be a single microstate and its time evolution to be unitary, such explanations were open to the well-known objection that, in a fully accurate description, the entropy must remain zero for all time.

Note that, conceptually, t above should presumably be thought of just as a ‘parameter’. Rather, it is tempting to identify the *time* at which a given unitary U ($= U(t)$ for some t , for some time-flow) *occurs*, with the entropy of $U\rho_0U^{-1}$. This remark seems to offer the prospect of at least a partial solution to the ‘problem of time in quantum gravity’.

Next, in order to remove an apparent obstacle to the validity of our thesis, we need to discuss in detail the nature of the quantum state of matter-gravity appropriate for the description of a black hole, as we now explain. Above, *we* have been taking it for granted that a black hole will always be correctly described by a single microstate of matter-gravity. *In the traditional picture* it is taken for granted, instead, that a black hole *cannot* be described by a single microstate. It is claimed, on the contrary, that it resembles a macrostate in that it needs to be described by a total density operator ρ with (total) non-zero (von Neumann) entropy, corresponding to a statistical mixture (i.e. a sum with suitable coefficients) of single microstates of the form $\rho_i = |\Psi_i\rangle\langle\Psi_i|$. As mentioned at the outset, this contradicts unitarity (in fact it’s not really clear what its physical interpretation could be) and this contradiction is known as the ‘information loss puzzle’⁶. To clarify this, we need to distinguish between:

- (a) the case of a black hole which is freshly formed from a collapsing mini-star modelled by

a state with zero matter-gravity entanglement in the distant past, and just beginning to evaporate, and

- (b) the stable equilibrium situation made possible by keeping the black hole in a suitably small box.

It seems that one of the main (at least implicit) reasons for the belief in the ‘information loss puzzle’ is that it was also taken for granted that, in case (b) – the perfect equilibrium situation the radiating black hole is, at the start, ‘striving’ to attain – the only physically possible equilibrium (total) states for this system are completely thermal. Instead, we shall now argue that the possible equilibrium (total) states of this system should equally include matter-gravity microstates in which the matter degrees of freedom and the gravitational degrees of freedom are separately in thermal equilibrium, but entangled. In other words that they should include single microstates which resemble ‘*double-KMS states*^{8,9}’ as we explain below. We suggest that this argument deserves to be considered as a resolution of the ‘information loss puzzle’ in that it removes the main reason for believing there to be such a puzzle.

To see that one expects there to be such single equilibrium microstates, we consider a simple analogy. Instead of matter and gravity, imagine a couplet of two real free scalar fields – called say the a-on field and the b-on field – in Minkowski space. Then, one knows that the generating functional (for all the n-point Wightman functions) of a thermal total state, at inverse temperature β , is given explicitly by:

$$\langle \exp(ia(f) + ib(g)) \rangle = \exp\left\{-\frac{1}{2}\langle Kf | \coth(\beta h/2) Kf \rangle - \frac{1}{2}\langle Kg | \coth(\beta h/2) Kg \rangle\right\}$$

where $a(f)$ and $b(g)$ are integrals over Minkowski space of the fields with test functions f and g , Kf and Kg denote the usual restrictions of the Fourier transforms of f and g to the future mass shell, h is the usual free field ‘one-particle Hamiltonian’, and the inner products in the exponent are defined by the usual Lorentz-invariant integral. However it is clear, from the work in refs. 8 and 9, that, aside from this total thermal (‘KMS’) state, one can also have the ‘double-KMS’ state

$$\begin{aligned} \langle \exp(ia(f) + ib(g)) \rangle = \exp\left\{-\frac{1}{2}\langle Kf | \coth(\beta h/2) Kf \rangle + \text{Re}\langle Kf | \text{cosech}(\beta h/2) Kg \rangle \right. \\ \left. - \frac{1}{2}\langle Kg | \coth(\beta h/2) Kg \rangle\right\} \end{aligned}$$

which has non-trivial correlations, i.e. entanglement, between the a-on field and the b-on field which are such that the total state is pure, i.e. a single microstate, but of course the partial states for the a-on field and the b-on field are identical to those of the thermal state and remain thermal. Note that this double-KMS state will have Green functions which are ‘periodic in imaginary time’, a property which is sometimes, mistakenly, believed to be exclusive to thermal states. Note also that one can give this model an amusing interpretation where the a-on field and the b-on field are identified with the restrictions of a single real scalar field to the left wedge and the right wedge of a fictitious Minkowski space, and the total (‘double KMS’) state is then analogous to the Minkowski vacuum

state. However, except in the two-dimensional massless case, where this identification can be achieved exactly by conformally mapping each of the wedges (in such a way that the wedge-preserving Lorentz boosts go over, in each case, into the forwards Minkowski time evolution), if one tried to take this identification literally, one would get⁸ a strange ‘one-particle Hamiltonian’.

Pursuing our analogy, *a-ons for matter, b-ons for gravity*, our suggestion, then, is that, also for a case (b) black hole, there will be a single microstate which resembles these double KMS states and this will be the physically relevant equilibrium state; *not* a totally thermal state. Concomitantly, we suggest that the full matter-gravity Euclidean path integral⁵ with the usual boundary conditions at the box and the usual periodicity in imaginary time is the generating functional for this single microstate, and that the traditional interpretation as a total matter-gravity thermal partition function is incorrect. Note however that, also with this new interpretation, if one takes the partial trace by integrating out all the matter fields, one will clearly still obtain a thermal partition function for the gravity degrees of freedom alone which coincides with that of refs. 4 and 5 to ‘zero-loops’. Hence we arrive at the conclusion that S_{gravity} is, to zero-loops, equal to (ii), as we assumed in our second argument for the validity of (2). We remark (A) If one were content to let our first argument for the validity of (2) stand alone, then one could turn that second argument into an *explanation* as to why (i) and (ii) approximately coincide. Moreover, since $S_{\text{matter}} = S_{\text{gravity}}$, we arrive at the *prediction* that the third entropy-like quantity (iii) the ‘entropy’ derived from the matter partition function obtained from the full matter-gravity Euclidean path integral by integrating out the gravity field

must also coincide, to a good approximation, with (i) and (ii). Prior to the present work, it was not clear that, or why, (i), (ii) and (iii) should have similar values and it was not clear which of (i), (ii), (iii), or possibly the sum of some of these, is the physical entropy of a black hole.

As further support for this suggestion, note that it would anyway seem unreasonable for the equilibrium state to which a radiating black hole is striving, to be a totally thermal state since, in such a state, matter and gravity would be unentangled and, as has often been explicitly or implicitly assumed, the matter and gravity partial von Neumann entropies would sum to the total von Neumann entropy. But, surely, because of interactions, the collapse process will tend to entangle an initially unentangled state leading to the failure of the von Neumann entropies to be additive. So just on these grounds, pure double-KMS-like states would seem better candidates for the physically relevant equilibrium states.

Finally, the assumptions we have made in this letter seem to tell us both *that* and *how* gravity must play a role in the mystery of the ‘collapse of the wave packet’: Our definition of entropy (2) may be interpreted as indicating that *the degrees of freedom of the gravitational field are unobservable*. We now propose to elevate this statement to a *principle*. With this principle, we may identify the algebra of observables with the algebra of operators A on the Hilbert space H_{matter} and, by (1), the expectation value of such an operator in the total microstate $\rho = |\Psi\rangle\langle\Psi|$ will be determined by the partial matter density operator ρ_{matter} through

$$\langle A \rangle = \text{trace}(\rho_{\text{matter}} A).$$

The physical entropy of ρ then becomes the von Neumann entropy of ρ_{matter} and the time evolution (3) between two times will act so as to transform a zero-entropy such partial density operator – corresponding to a total initial microstate which is unentangled – into a density operator with non-zero, and by our arguments above, ever increasing, entropy. Thus, with our principle, as far as the (matter) observables are concerned, the gravitational field must play a role similar to that which would be played, in the traditional view, by a combination of a unitary time-evolution and a continual process of ‘measurement’, i.e. the ‘reduction’ part of measurement; perhaps the ‘collapse’ remains mysterious. Previously there had been attempts to understand such ‘decoherence’ in terms of ‘the environment’ but they did not really work because, as long as the environment consisted of degrees of freedom which were observable in principle, the wave function could, at best, be reduced ‘for all practical purposes’, but never *actually* be reduced. Instead, by constituting an environment which is in principle unobservable, the gravitational field gives us an *absolute notion of reduction* thus finally offering at least a partial explanation for why we don’t see ‘Schrödinger cats’ etc. In the ‘measurements’ which are taking place in the decoherence process, the ‘cut’⁷ between the unitary time evolution and the ‘reduction’ is now understood not to occur, as von Neumann argued, at some arbitrarily located stage, between ‘system’ and ‘observer’, but rather to be located, precisely, where the matter ‘open system’ makes contact with the ‘gravity environment’. All this clarifies the foundations of quantum mechanics in much the same way as our objective notion of coarse graining clarified the foundations of statistical mechanics. In fact, with the above principle, we have achieved not only a clarification, but also a *unification* of the two subjects; *entropy increase and decoherence are seen to be two aspects of the same thing*.

Hawking’s calculation¹ of black hole evaporation, twenty four years ago, provided a crucial clue which has enabled us, here, to achieve a new understanding of thermodynamics and to find a single explanation for the origin both of entropy increase and of decoherence. Contradicting Einstein, Hawking⁶ said that ‘God not only plays dice, but sometimes throws the dice where we cannot see them’. Here, we have confirmed that this is true, but have argued that the dice we cannot see are nothing other than the degrees of freedom of the gravitational field itself; while remaining hidden, they ensure the overall unitarity of the theory.

The framework we have arrived at is remarkably close to conventional quantum mechanics. But it has some important new features: unobservable degrees of freedom, and a built-in irreversibility; time has a beginning, and the initial state is unentangled. The challenge it poses is to construct a full quantum theory of gravity within this new paradigm.

1. Hawking, S.W. Black hole explosions? *Nature* **248**, 30-31 (1974).
2. Hawking, S.W. Particle creation by black holes. *Commun. Math. Phys.* **43**, 199-220, (1975).
3. Wald, R.M. Black holes and thermodynamics. in *Proceedings of the Chandra Symposium: “Black Holes and Relativistic Stars”* (ed. Wald, R.M.) (U. of Chicago Press, to appear) (gr-qc/9702022)
4. Gibbons, G.W. & Hawking, S.W. Action integrals and partition functions in quantum gravity. *Phys. Rev.* **D15**, 2752-2756 (1977).

5. Hawking, S.W. The path integral approach to quantum gravity. in *General Relativity, An Einstein Centenary Survey* (eds Hawking, S.W., Israel, W.) 746-785 (Cambridge University Press, 1979).
6. Hawking, S.W. Breakdown of predictability in gravitational collapse. *Phys. Rev. D* **14**, 2460-2473 (1976).
7. von Neumann, J. *The Mathematical Foundations of Quantum Mechanics* (Princeton University Press, 1955).
8. Kay, B.S. The double wedge algebra for quantum fields on Schwarzschild and Minkowski spacetimes. *Commun. Math. Phys.* **100**, 57-81, (1985).
9. Kay, B.S. Purification of KMS states. *Helvetica Physica Acta* **58**, 1030-1040 (1985).

Acknowledgements. I am very grateful to Leslie Sage for reading, and constructively criticizing, a first version of this work and to Wanda Andreoni and Alessandro Curioni who (during the week 15-22 January 1998) did the same for successive further versions. Especially, I am indebted to Alessandro Curioni for reminding me that entropy is time's arrow and for insisting that, of the two concepts, entropy is the more fundamental.

Correspondence should be addressed to the author (e-mail bsk2@york.ac.uk).